

Signatures and alternating tangle decompositions

Adam Lowrance

Joint with Cody Armond and Oliver Dasbach


Joint Mathematical Meetings

January 8, 2016




Turaev surfaces


The **A**-resolution and **B**-resolution of a crossing:



Create Turaev surface from link diagram:



Turaev surface: $P(3,3,-2)$




Turaev surfaces


- ❖ Start with cobordism between all-**A** and all-**B** state.
- ❖ Cap off boundary components with disks.
- ❖ Turaev genus $g_T(L)$: Minimal genus among all Turaev surfaces.
- ❖ The link projection on the Turaev surface is alternating.
- ❖ $g_T(L) = 0$ if and only if the link is alternating.

Turaev surfaces


- ❖ Replace crossings with disks.
- ❖ Replace edges with possibly twisted bands.
- ❖ Cap off boundary components with disks.



Turaev surfaces




Turaev surfaces




Alternating decompositions


- ❖ Separate a link diagram into its maximal alternating regions.
- ❖ Mark each non-alternating edge with two points.
- ❖ Inside each face, connect the marked points with arcs as below.




Alternating decompositions



Alternating decompositions



Alternating decompositions




Alternating decompositions

- ❖ The alternating decomposition determines a graph: curves \rightarrow vertices; non-alternating arcs \rightarrow edges.
- ❖ The graph determines the genus of the Turaev surface.
- ❖ There is a recursive algorithm on graphs to compute the genus of the Turaev surface.

Turaev genus one diagrams

(Armond, L.; Kim) Let D a prime link diagram. The Turaev surface of D is genus one if and only if the alternating decomposition graph is a doubled cycle of even length.



Classification results

- ❖ Lengthening/shortening doubled paths preserves the genus of the Turaev surface.
- ❖ There are a finite number of double-path equivalence classes of alternating decomposition graphs of any fixed Turaev genus.
- ❖ There are five equivalence classes for Turaev genus two.

Determinant and *Signature* of a knot

V Seifert matrix for knot

Determinant:

$$|\det(V + V^T)| = |\Delta_K(-1)| = |V_K(-1)|$$

Signature:

$$\sigma(V + V^T)$$

Determinant and Signature of a knot

Determinant of a knot is odd and
the signature is even

Known: Changing a positive to
a negative crossing:

$$\sigma(K_-) - 2 \leq \sigma(K_+) \leq \sigma(K_-)$$

Signature for Turaev genus 0 links (alternating links)

(Gordon-Litherland, Traczyk) For an
alternating knot K with alternating
diagram D

$$\sigma(K) = s_A(D) - c_+(D) - 1 = -s_B(D) + c_-(D) + 1$$

Signature and Turaev genus


The genus $g_T(D)$ of the Turaev surface is given by

$$\begin{aligned} 2g_T(D) &= 2 + c(D) - s_A(D) - s_B(D) \\ &= (-s_B(D) + c_-(D) + 1) - (s_A(D) - c_+(D) - 1) \end{aligned}$$

(Dasbach, L.) For a knot K with diagram D

$$s_A(D) - c_+(D) - 1 \leq \sigma(K) \leq -s_B(D) + c_-(D) + 1$$

Example: Signature of $P(3,3,-2)$



$$s_A(D) - c_+(D) - 1 = 3 - 8 - 1 = -6$$

$$g_T(D) = 1$$

$$-6 \leq \sigma(K) \leq -4$$

Signature and Turaev genus

(Murasugi) For a knot K

$$\sigma(K) = \det(K) - 1 \pmod{4}.$$


(Dasbach, L.) For a knot K with diagram D of Turaev genus 1 the signature is determined by:

$$\sigma(K) = s_A(D) - c_+(D) \pm 1$$

and

$$\sigma(K) = \det(K) - 1 \pmod{4}.$$

Example: Signature of $P(3,3,-2)$





$$\sigma(K) = -6 \text{ or } \sigma(K) = -4$$

$$\det(K) = 3$$


$$\sigma(K) \equiv 2 \pmod{4}$$

$$\Rightarrow \sigma(K) = -6$$

Closures of tangles




$N(R)$



$D(R)$

Signature and Turaev genus



(Dasbach, L.) Let K be a knot with diagram as above. Then

$$\sigma(K) = \pm 1 + \sum \sigma(N(R_i)),$$

or

$$\sigma(K) = \pm 1 + \sum \sigma(D(R_i)).$$

Signatures of Montesinos knots

- ❖ Murasugi computes the signature of the rational knot $K(p/q)$ from p and q .
- ❖ Qazaqzeh, Yasein, and Abu-Qamar compute the signature of $K(p/q)$ from the continued fraction expansion of p/q .
- ❖ Champanerkar and Ording give a formula for the determinant of a Montesinos link.
- ❖ The theorem on the previous slide then tells us how to compute the signature of a Montesinos knot.

Thank you